

# Directional Dark Matter Search and Velocity Distribution




Keiko I. Nagao (National Institute of Technology, Niihama College, Japan)

Collaboration with

Tatsuhiko Naka (Nagoya Univ.) and Mihoko Nojiri (KEK & IPMU)

# Dark Matter


## Bullet cluster





## Weak Lensing

# Evidences

## Galactic Rotation Curve




## CMS




# Constraints for DM


## Direct Detection




## Collider (LHC)



## Relic Abundance



## Indirect Detection





# Direct Detection

## □ Direct search for DM

- Detecting **the recoil energy** that a DM particle scatters a target nucleus.
- CDMS, XENON, LUX, ...

## □ Constraint for DM-nucleon interaction cross section can be obtained from the event number

$$R = \underbrace{N_T n_\chi}_{\text{Experiment}} \int_{E_{R,\min}} dE_R \underbrace{\int_{v_{\min}}^{v_{\max}} d^3v f(v)}_{\text{Astronomy}} \underbrace{\frac{\tilde{\sigma} A m_A}{2v\mu_A^2}}_{\text{Particle + nuclear phys.}}$$



# OUTLINE

1. ~~Introduction~~
2. Directional Dark Matter Detection
3. Velocity Distribution of Dark Matter
4. Nuclear Emulsion Detector
5. Velocity Distribution Observed in the Directional Detector

# To next generation: Directional detection

## ■ Directional Detection

- detecting not only the recoil energy but also **direction where DM comes from**.

## ■ Advantages

### - **Powerful back ground rejection**

BG is isotropic, on the other hand DM signal is expected to come from the direction of the cygnus.



### - **Annual Modulation**

Direction of DM wind toward the Earth seasonally changes.


### - **Daily oscillation**

The Earth's rotation can also changes the number of DM signals.


DM wind



# Directional Dark Matter Searches



# Directional Searches





# We can be more ambitious?

**Dark  
Matter**

**Direction**

**Detection**



# We can be more ambitious?



# Being more ambitious...


- Velocity distribution  
In the directional DM search, **it can be possible to make a constraint for  $f(\mathbf{v})$ .**
- Constraint from direct detections depends on DM distribution

The diagram shows the equation for the direct detection rate  $R$ . The terms are annotated as follows:

- $N_T$  and  $n_\chi$  are enclosed in a dashed orange box labeled "Experiment".
- The energy integral  $\int_{E_{R,\min}} dE_R$  is enclosed in a dashed orange box labeled "Experiment".
- The velocity integral  $\int_{v_{\min}}^{v_{\max}} d^3v f(v)$  is enclosed in a solid blue box labeled "Astronomy". A yellow arrow points down to this box.
- The cross-section term  $\frac{\tilde{\sigma}_{AM_A}}{2v\mu_A^2}$  is enclosed in a solid red circle labeled "Particle + nuclear phys."

- We should know correct DM distribution to derive appropriate constraints for the interaction.

# DM Velocity Distribution -Standard Distribution-



Catena and Ullio (2012)

## ■ Maxwell distribution

$$f(v) = \frac{1}{(\pi v_0^2)^{3/2}} e^{-(v+v_E)^2/v_0^2}$$



$v_0$ : velocity of the Solar system

$v_E$ : Earth's velocity relative to DM

## ■ Is DM distribution surely this kind of shape?

# Debris Flow




- N-body simulation in which subhalos falling into the Milky Way



Kuhlen, Lisanti & Spergel (2012)

# Co-rotating DM



- N-body simulation including baryons and gas
  - DM co-rotates with baryons in the galaxy.
  - Anisotropic distribution



Ling, Nezri, Athanassoula & Teyssier (2009)

# Distribution changes the constraint for interaction

- Applying non-standard distribution (with other factors, like isospin violating, inelastic scattering...) can improve the situation to explain the discrepancy between positive and negative results of direct searches.



Cline, Liu, Xue (2012)

# Nuclear Emulsion Detector




# Nuclear Emulsion

## ■ Nuclear Emulsion



- A kind of photographic film
- 3D tracking detector for charged particle:

Charged particle can expose silver halide crystals (AgBr) in films. After development treatment, the track appears as silver grains.



# Concept of DM detection with nuclear emulsion



- Detection of **recoiled nucleus** from DM-nucleon scattering



Rotation of the Earth can change the direction of detector toward the DM wind, however, detector direction can be kept to be against DM wind by an equatorial telescope.

# Nuclear Emulsion Detector (I)

- Underground facility which had been used for OPERA project
- In research & development
- Taking BG data




**GROUP:**

- Nagoya University
- Napoli university
- Padova university
- LNGS

# Nuclear Emulsion Detector (II)

## ■ Advantages

- High sensitivity :  
solid target + large mass  
(O(100) kg)
- High spatial resolution  
Angular resolution: 15-20°  
Spatial resolution: 100 nm
- Low cost (150,000yen/kg~870£/kg)




# Energy threshold

## ■ Target


- Ag, Br, C, N, O
- Energy threshold : depends on target  
(~33 keV for C, N, O and  
~150 keV for Ag, Br)

## ■ For O(10)-O(1000)GeV mass DM

- Typical recoil energy : O(1)-O(100)keV
- Required resolution is submicron  
(~O(100)nm) track length



## Weight Ratio




# **Velocity Distribution observed in the Directional Detector**

# Can we distinguish the velocity distribution?

- In the directional direct search, we can see both the scattering angle and the recoil energy.



Calculation





- Monte Carlo simulation
- Simple elastic scattering
- Scattering angle– Recoil energy (track length) distribution

# Track Length for light DM

- DM mass :200 GeV
- Target: realistic component of emulsion (Ag, Br, C, N, O)





$\cos(\theta)$  - track length [nm]







# Track Length for heavy DM

- DM mass :200 GeV
- Target: realistic component of emulsion (Ag, Br, C, N, O)



cos( $\theta$ ) - track length [nm]



# Summary & Discussion

- I discussed the possibility to distinguish the distribution models of dark matter in the direct detection, focusing on the nuclear emulsion experiments.
- Distribution shape of the scattering angle and the energy density is affected by the distribution model. If the number of DM signal is enough, it seems possible to give a constraint for the velocity distribution model.

# Backup Slides

# BG rejection -summary-

- Radioactive sources from outside :  $\beta$ ,  $p$ ,  $\mu$ 
  - Sensitivity control, point-like signal
- Internal BG sources :  $\beta$ , ( $\gamma$ )
  - $^{40}\text{K}$  mixed in when  $\text{KBr} \rightarrow \text{AgBr}$ , can be avoided by using  $\text{NaBr}$  instead of  $\text{KBr}$
  - $^{14}\text{C}$  ( $\beta$ -ray induced by  $\gamma$  makes the grains which has Plasmon resonance effects, i.e., we can distinguish them by color obs.)
- Neutron from rocks
  - Neutron shield, sensitivity control
- Others
  - Underground, isotropic angular distribution

# Contents of nuclear emulsion

		Weight(%)	$A_i$ (abundance)	
AgBr crystal	Ag	39.65	107(51.84)	109(48.16)
	Br	29.01	79(50.69)	81(49.31)
gelatin	O	11.76	16	
	C	11.72	12(98.9)	13(1.1)
	N	4.57	14	
	H	2.27	1	
	S	0.05	32(95.02)	34(4.2)
	I	0.96	127	

# Periodic Table

**Periodic Table of the Elements**

© www.elementsdatabase.com

1 H																	2 He														
3 Li	4 Be											5 B	6 C	7 N	8 O	9 F	10 Ne														
11 Na	12 Mg											13 Al	14 Si	15 P	16 S	17 Cl	18 Ar														
19 K	20 Ca	21 Sc	22 Ti	23 V	24 Cr	25 Mn	26 Fe	27 Co	28 Ni	29 Cu	30 Zn	31 Ga	32 Ge	33 As	34 Se	35 Br	36 Kr														
37 Rb	38 Sr	39 Y	40 Zr	41 Nb	42 Mo	43 Tc	44 Ru	45 Rh	46 Pd	47 Ag	48 Cd	49 In	50 Sn	51 Sb	52 Te	53 I	54 Xe														
55 Cs	56 Ba	57 La	72 Hf	73 Ta	74 W	75 Re	76 Os	77 Ir	78 Pt	79 Au	80 Hg	81 Tl	82 Pb	83 Bi	84 Po	85 At	86 Rn														
87 Fr	88 Ra	89 Ac	104 Unq	105 Unp	106 Unh	107 Uns	108 Uno	109 Une	110 Unn																						
																		58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
																		90 Th	91 Pa	92 U	93 Np	94 Pu	95 Am	96 Cm	97 Bk	98 Cf	99 Es	100 Fm	101 Md	102 No	103 Lr